
No.



9000265

THE UNIVERD SHAYIES OF ANTERIOA

TO ALL TO WHOM THESE PRESENTS SHALL COME;

Northrup King Co.

Tellicreas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF eighteen years from the date of this grant, subject to the payment of the required fees and periodic replenishment of viable basic seed of the variety in a public repository as provided by LAW, the right to exclude others from selling the variety, or offering it for sale, or reproducing it, importing it, or exporting it, or using it in producing a hybrid or different tety therefrom, to the extent provided by the Plant Variety Protection Act

AT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

SOYBEAN

'S64-23'

In Testimony Watercot, I have hereunto set my hand and caused the seal of the Plant Variety Protection Office to be affixed at the City of Washington, D.C.

this 31st day of August in the year of our Lord one thousand nine hundred and ninety-two.

Secretary of Agriculture

Attosk

Kenneth Hon

Plant Variety Protection Office Agricultural Marketing Service Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W. Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

FORM APPROVED: OMB 0581-0055, Exprise 1/31/91

U.S. DEPARTMENT OF A AGRICULTURAL MARKET APPLICATION FOR PLANT VARIET (Instructions on I	ING SERVICE Y PROTECTIO	N CERTIFICATE	d c	Application is required in order to determine if a plant variety protection entificate is to be issued (7 U.S.C. 2421). Information is held confidential until entificate is issued (7 U.S.C. 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)		2. TEMPORARY DESIGNA EXPERIMENTAL NO.	ATION OR 3.	VARIETY NAME
Northrup King Co.		X8964		S64-23
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area	ode)	FOR OFFICIAL USE ONLY
P. O. Box 959			P'	VPO NUMBER
Minneapolis, MN 55440		612-593-73	33	9000265
				F Date . 6, 1990
6. GENUS AND SPECIES NAME	7. FAMILY NAME (Botai	nical)		Time
Glycine max	Leguminos	ae		N
8: CROP KIND NAME (Common Name)	<u> </u>	-		F Filing and Examination Fee:
	9	DATE OF DETERMINATION		E 32150.
Soybean		March, 1987	ļ. ·	S Date
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGAN Corporation	IIZATION (Corporation, pe	artnership, association, etc.)		Sept. 6,1990 C Certificate Fee:
		·		: 250,00
11. IF INCORPORATED, GIVE STATE OF INCORPORATION Delaware	12. [DATE OF INCORPORATION 1976		E Cug 12,1992
Robert W. Romig Northrup King Co. P. O. Box 959 Minneapolis, MN 55440 14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Folia). a. X Exhibit A, Origin and Breeding History of the Variety b. X Exhibit B, Novelty Statement. c. X Exhibit C, Objective Description of Variety. d. Exhibit D, Additional Description of Variety. e. X Exhibit E, Statement of the Basis of Applicant's Ownershit f. X Seed Sample (2,500 viable untreated seeds). Date Seed g. X Filing and Examination Fee (\$2,150) made payable to "T 15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOME Protection Act.) YES (II "YES." answer items 16 and 17 bells of the Company of the Compan	Sample mailed to Plant reasurer of the United in the Unite	erse) I Variety Protection Office States." ILY AS A CLASS OF CERTIFIE "NO," skip to item 18 below) TO ITEM 16, WHICH CLASSES DUNDATION		ON BEYOND BREEDER SEED?
YES (II "YES," give names of countries and dates) X NO 20. The applicant(s) declare(s) that a viable sample of basic serequest in accordance with such regulations as may be appl	icable.			
The undersigned applicant(s) is (are) the owner(s) of this uniform, and stable as required in section 41, and is entitled. Applicant(s) is (are) informed that false representation here.	d to protection under	the provisions of section	42 of the Pla) that the variety is distinct, int Variety Protection Act.
	CAPACITY OF	-		DATE
SIGNATURE OF APPLICANT (Owner(s))	-	President, Rese	arch	8/31/90
SIGNATURE OF ARRUSTANT (Ourself 1)	CAPACITY OF	D TITLE		DATE
SIGNATURE OF APPLICANT [Owner(s)]	CAPACITY OF			

EXHIBIT A

Origin and Breeding History of S64-23

In 1976, the Coker's Pedigreed Seed Company group at Hartsville, SC made the cross J77-46 x (Coker 136*3 x Centennial)F1 from which the variety S64-23 is derived. J77-46 was later released as the variety, Bedford.

The F1 was advanced to F2 in the field at Hartsville in 1977, and the F2 was grown in the field the following year and harvested in bulk. The F3 was grown over winter in the greenhouse during 1978-79 and advanced to F4 using single seed descent. The F4 was grown in the field in 1979 and single plant selections were made at The F5 progeny from these plants were evaluated for resistance to Race 3 of soybean cyst nematode during the winter of Resistant F5 lines were grown in progeny rows in the field during 1980 and PR #653 was selected and harvested in bulk, and was one of a number of selected progeny rows which were screened as resistant to Race 4 of cyst nematode during the winter of 1980-81. F6 rows were grown at Richland, Indiana during 1981 as part of a "Cyst Nematode Advanced Progeny" nursery and progeny #CNAP118 was selected and harvested in bulk. A further F7 generation of progenies were evaluated at Hartsville during 1982 and row #4329 was selected and again harvested in bulk. point the material was classified as early group VI and was moved to the mid-South program then located at West Memphis, Arkansas and was designated Co 83M-32. From 1983-84, Co 83M-32 was evaluated in yield trials throughout the mid-southern United States.

During this period, the line was characterized as having white flowers, tawny pubescence, tan pods, and seeds with a black hilum and absence of seed coat luster. Co 83M-32 was further evaluated in advanced yield trials across a wide range of environments from 1985-89 when, in addition to establishing its yield superiority, it was also found to be resistant to frogeye leafspot and southern root knot nematode. As a result it was decided to release the line as \$64-23 in 1989.

Breeder seed was produced in 1988 by bulking seed from similar progeny rows grown and harvested separately over two generations. Foundation seed was produced and approved by the Arkansas State Plant Board in 1989. Varietal purity will be maintained using progeny rows and roguing as needed.

S64-23 is a uniform, stable variety. Occasional off-type plants with purple flowers, tawny pubescence, and brown hilum and a further off-type with white flowers, gray pubescence and buff hilum have been observed. These were assumed to have resulted from mixture. During the seven years of testing and three years of seed increase, we have observed no other off-types except for minor environmentally induced variations in the intensity of hilum pigmentation.

2

EXHIBIT B (AMENDED)

Novelty Statement for Soybean Variety '864-23'

Soybean variety 'S64-23' is most similar to the variety Bedford. It can be differentiated from Bedford on the basis of reactions to Southern root knot nematode (Melo-idogyne incognita) and to Frogeye leaf spot (Cercospora sojinae). Variety 'S64-23' is resistant to Southern root knot nematode whereas Bedford is susceptible. Variety 'S64-23' also has greater resistance to Race 4 of Frogeye leaf spot, as shown in the table below.

Table 1. Average leaf area affected from challenge by spores of *Cercospora sojinae*

Variety	Race 4, Isolate			
7422003	CS-03	CS-05		
	8	8		
'S64-23'	3.9	10.6		
Bedford	16.6	25.0		
Blackhawk (susceptible)	39.4	43.6		
Davis (resistant)	0.0	0.0		
LSD _{.05}	8.5	13.5		

This Frogeye leaf spot test was conducted in a greenhouse at Auburn University in the spring of 1992. Plants were inoculated at the VI-V2 leaf stage with a suspension of spores in water with Tween 20 surfactant at a concentration of from 4x10⁴ to 6x10⁴ spores/ml. Inoculated pots were placed on a greenhouse bench in a randomized complete block design with four replications (four pots). Scores represent the leaf area affected, as percent, from three leaflets/plant with three plants/pot.

EXHIBIT C (Soybean)

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK, MEAT, GRAIN & SEED DIVISION PLANT VARIETY PROTECTION OFFICE BELTSVILLE, MARYLAND 20705

OBJECTIVE DESCRIPTION OF VARIETY SOYBEAN (Glycine max L.)

	- Toryeme max E.	<u> </u>
NAME OF APPLICANT(S)	TEMPORARY DESIGNATION	VARIETY NAME
Northrup King Co.	X8964	S64-23
ADDRESS (Street and No., or R.F.D. No., City, State, and Zip Cod	(e)	FOR OFFICIAL USE ONLY
P. O. Box 959		PVPO NUMBER
Minneapolis, MN 55440	i	0000045
Attention: R. W. Romig		9000265
Choose the appropriate response which characterizes the var	riety in the features described l	pelow. When the number of significant digits
in your answer is fewer than the number of boxes provided,	place a zero in the first box w	hen number is 9 or less (e.g., 0 9).
1. SEED SHAPE:		
	.	•
L W	[T]	
1 = Spherical (L/W, L/T, and T/W ratios = < 1.2)	2 = Spherical Flattened	(L/W ratio > 1.2; L/T ratio = < 1.2)
3 = Elongate (L/T ratio > 1.2; T/W = < 1.2)	4 = Elongate Flattened (L/T ratio > 1.2; T/W > 1.2)
2. SEED COAT COLOR: (Mature Seed)		
1 4 - Valley 0 - Court	4 - Black	/P:£.1
1 = Yellow 2 = Green 3 = Brown	4 = Black 5 = Other	Specify)
3. SEED COAT LUSTER: (Mature Hand Shelled Seed)		
		•
1 = Dull ('Corsoy 79'; 'Braxton') 2 = Shiny ('Nebs	oy'; 'Gasoy 17'}	
4. SEED SIZE: (Mature Seed)		
		•
1 3 Grams per 100 seeds	•	
5. HILUM COLOR: (Mature Seed)		
5. HILUM COLOR: (Mature Seed)		
6 1 = Buff 2 = Yellow 3 = Brown	4 = Gray 5 = Imperfect Bis	ck 6 = Black 7 = Other (Specify)
6. COTYLEDON COLOR: (Mature Seed)		
1 1 = Yellow 2 = Green	•	
7. SEED PROTEIN PEROXIDASE ACTIVITY:		
1 1 = Low 2 = High		
1 1 2 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		
8. SEED PROTEIN ELECTROPHORETIC BAND:		
	•	
2 1 = Type A (SP1 ^a) 2 = Type B (SP1 ^b)		
9. HYPOCOTYL COLOR:		
s, strocorrection.		
1 2 1	h bronze band below cotyledons (*	Woodworth'; 'Tracy')
3 = Light Purple below cotyledons ('Beeson'; 'Pickett 71') 4 = Dark Purple extending to unifoliate leaves ('Hodgson';	'Coker Hampton 266A')	
	·	
10. LEAFLET SHAPE:	•	
3 1 = Lanceolate 2 = Oval 3 = Ovate	4 * Other (Specify)	
		<i>,</i> /-

FORM LMGS-470-57 (2-82)

Page 2 of 4

4.5.	LEAF	LET SIZE:		
	2	1 = Small ('Amsoy 71'; 'A5312') 3 = Large ('Crawford'; 'Tracy')	2 = Medium ('Corsoy 79'; 'Gasoy 17')	· · · · · · · · · · · · · · · · · · ·
12.	LEAF	COLOR:		
	2	1 = Light Green ('Weber'; 'York') 3 = Dark Green ('Gnome'; 'Tracy')	2 = Medium Green ('Corsoy 79'; 'Braxton')	
13.	FLOW	ER COLOR:		
	1	1 = White 2 = Purple	3 = White with purple throat	
14.	POD C	OLOR:		
	1	1 = Tan 2 = Brown	3 = Black	
15.	PLAN	F PUBESCENCE COLOR:	:	
	2	1 = Gray 2 = Brown (Tawny)		
16.	PLANT	T TYPES:		
	2	1 = Slender ('Essex'; 'Amsoy 71') 3 = Bushy ('Gnome'; 'Govan')	2 = Intermediate ('Amcor'; 'Braxton')	
17.	PLANI	T HABIT:		
11	1	1 = Determinate ('Gnome'; 'Braxton') 3 = Indeterminate ('Nebsoy'; 'Improved Potential Pot	2 = Semi-Determinate ('Will') elican')	
18.	MATU	RITY GROUP:		
	9		The second secon	7 = IV 8 = V
		1 = 000 2 = 00 3 = 0 9 = VI 10 = VII 11 = VI	4 = I 5 = II 6 = III II 12 = IX 13 = X	7-1V 8-V
19.	DISEA		II 12 = IX 13 = X	/-IV
19.		9 = VI	II 12 = IX 13 = X	/-IV
19.	BACT	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES:	II 12 = IX 13 = X Susceptible; 2 = Resistant)	/-IV
19.		9 = VI	II 12 = IX 13 = X Susceptible; 2 = Resistant)	/-IV
19.	BACT	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES:	II 12 = IX 13 = X Susceptible; 2 = Resistant)	/-IV
19.	BACT	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli v	II 12 = IX 13 = X Susceptible; 2 = Resistant)	/-IV
	2 1 2	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea)	Il 12 = IX 13 = X Susceptible; 2 = Resistant) var. sojensis)	,-IV
	2 1 2	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci)	Il 12 = IX 13 = X Susceptible; 2 = Resistant) var. sojensis)	,-IV
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli v Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES:	Il 12 = IX 13 = X Susceptible; 2 = Resistant) var. sojensis)	
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli v Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina)	Susceptible; 2 = Resistant) var. sojensis) RWS - 1/10/92	2 Other (Specify)
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2	Susceptible; 2 = Resistant) var. sojensis)	2 Other (Specify) Resistant to common
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2 R	Susceptible; 2 = Resistant) var. sojensis) Race 3	2 Other (Specify)
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2	Susceptible; 2 = Resistant) var. sojensis) Race 3	2 Other (Specify) Resistant to common
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2 R	Susceptible; 2 = Resistant) var. sojensis) Race 3	2 Other (Specify) Resistant to common
	BACT 2 1 2. FUNGA	9 = VI 10 = VII 11 = VI SE REACTION: (Enter 0 = Not Tested; 1 = TERIAL DISEASES: Bacterial Pustule (Xanthomonas phaseoli vi Bacterial Blight (Pseudomonas glycinea) Wildfire (Pseudomonas tabaci) AL DISEASES: Brown Spot (Septoria glycines) Frogeye Leaf Spot (Cercospora sojina) Race 1 Race 2 R Target Spot (Corynespora cassiicola) Downy Mildew (Peronospora trifoliorum vi	Susceptible; 2 = Resistant) var. sojensis) Race 3 2 Race 4 Race 5	2 Other (Specify) Resistant to common

FORM LMGS-470-57 (2-82)

19, DISE	ASE REACTION:	(Enter 0 = Not Tested; 1 = Susceptible; 2 = R	esistant) (Continued)	·
FL	INGAL DISEASES	: (Continued)		
•	Pod and Stem	Blight (Diaporthe phaseolorum var; sojae)		
	Purple Seed St	ain (Cercospora kikuchii)	•	•
	Rhizoctonia R	oot Rot (Rhizoctonia solani)		
· •	Phytophthora	Rot (Phytophthora megasperma var. sojae)	•	
1	Race 1	1 Race 2 1 Race 3 1	Race 4 1 Race 5	1 Race 6 1 Race 7
1	Race 8	1 Race 9 Other (Specify)		
VI	ப் RAL DISEASES:			
	7	obacco Ringspot Virus)		
	- -	(Bean Yellow Mosaic Virus)	•	
F	1	c (Cowpea Chlorotic Virus)		
	i			
_	i	ean Pod Mottle Virus)		
L		loybean Mosaic Virus)	÷	
NE	MATODE DISEA			
- ا	Soybean Cyst	Nematode (Heterodera glycines)		
<u> </u>	_ Race 1 <u> </u>	Race 2 2 Race 3 2	Race 4 (14) Other (S	pecify)
	Lance Nemato	de (Hopiolaimus Colombus)	•	
2	Southern Roo	t Knot Nematode (Melaidogyne incognita)		
	Northern Room	t Knot Nematode (Meloidogyne Hapla)		
1	Peanut Root K	not Nematode (Meloidogyne arenaria)		
	Reniform Nem	atode (Rotylenchulus reniformis)		•
F	OTHER DISE.	ASE NOT ON FORM (Specify):	<u> </u>	
20. PHYS	SIOLOGICAL RES	PONSES: (Enter 0 = Not Tested; 1 = Suscept	ible; 2 = Resistant)	
		on Calcareous Soil		
· L	Other (Specify	/		
21. INSE	CT REACTION:	Enter 0 = Not Tested; 1 = Susceptible; 2 = Re	sistant)	
	Mexican Bean	Beetle (Epilachna varivestis)	·	
	Potato Leaf Ho	opper (Empoasca fabae)		
	Other (Specify)		
22. INDI	CATE WHICH VA	RIETY MOST CLOSELY RESEMBLES THAT	T SUBMITTED.	· · · · · · · · · · · · · · · · · · ·
СН	ARACTER	NAME OF VARIETY	CHARACTER	NAME OF VARIETY
Plant	Shape	Bedford	Seed Coat Luster	Coker 136
Leaf S	Shape	Centennial	Seed Size	Bedford
Leaf (Color	Stonewall	Seed Shape	Coker 136
Leaf S	Size	Centennial	Seedling Pigmentation	Coker 136
			·	4

23. GIVE DATA FOR SUBMITTED AND SIMILAR STANDARD VARIETY: Paired Comparison Data

VARIETY	NO. OF DAYS MATURITY		CM PLANT HEIGHT	LEAFLET SIZE		SEED CONTENT		SEED SIZE	NO.
				CM Width	CM Length	% Protein	% Oil	G/100 SEEDS	SEEDS/ POD
S64-23 Submitted	154	2.2	102	8.5	13.3	35.2	16.4	13.4	2-3
RA606 Name of Similar Variety	156	2.2	99	5.9	11.5	35.1	16.7	12.2	2-3

PUBLICATIONS USEFUL AS REFERENCE AIDS FOR COMPLETING THIS FORM:

- 1. Caldwell, B.E., ed. 1973. Soybeans: Improvement, Production, and Uses. Amer. Soc. Agron. Monograph No. 16.
- 2. Buttery, B.R. and R.I. Buzzell. 1968. Peroxidase activity in seeds of soybean varieties. Crop Sci., 8: 722-725.
- 3. Hymowitz, T. 1973. Electrophoretic analysis of SBTI-A₂ in the USDA soybean germplasm collection. Crop Sci., 13: 420-421.
- 4. Payne, R.C. and L.F. Morris. 1976. Differentiation of soybean cultivars by seedling pigmentation patterns. J. Seed Technol. 1: 1-19.

EXHIBIT E

Statement of the Basis of Applicant's Ownership

Soybean variety S64-23 was developed by the Northrup King Co. soybean breeding staff from germplasm sources cited in Exhibit A of this application. Northrup King Co. believes that the variety is novel as defined in the Plant Variety Protection Act and, therefore, that Northrup King Co. is the sole owner of the variety.